



Studies on Prevalence of Platyhelminth Parasites of freshwater fishes from Sukhana Dam at Aurangabad District (MS) India

Nimbalkar R K* and Deolalikar RV

Department of Zoology, Vinayakrao Patil Mahavidyalaya, Vaijapur, District Aurangabad (M.S.)

*rknimbalkar@gmail.com

Article Info

Received: 28-05-2015,

Revised: 26-06-2015,

Accepted: 30-06-2015

Keywords:

Cestode, trematode, Sukhana Dam, population dynamics.

Abstract

In present study at Sukhana dam, we collected 147 fish species of *Wallago attu* ($n=22$), *Mastacembalus armatus* ($n=95$) and *Clarius batrachus* ($n=30$). Out of these, 101 fishes were found infected with two genus of cestode parasites viz., *Senga sukhaensis* and *Gangesia attuensis* we also collected 10 species of trematode parasites *Phyllodistomum batrachusi* from urinary bladder of fresh water cat fish *Clarius batrachus* during January 2010 to December 2012. The population dynamics shows the prevalence, mean intensity, abundance and dominance of the collected cestode and trematode parasites. During this study we also observed protozoans, crustaceans, Monogenea as well as Acanthocephala and Nematodes.

INTRODUCTION

India is among the 17 mega diversity countries (Mittermeier *et al.*, 1997) and hosts as many as 55 families of freshwater fish (Froese and Pauly, 2015). For the last few decades, fishes have been extensively used as food for human consumption in the Indian subcontinent and thus contribute substantially to its economy. In India it is estimated that about 10 million tons of fish are required to meet the annual demand of fish proteins as compared to an actual annual production of only 3.5 million tons (Shukla and Upadhyay, 1998). Catfishes are an important in fish fauna of wetlands and many of them are economically important as a food source of high nutritive value. Schimidt and Roberts, (2000) have reported that endoparasitic helminths, with indirect life cycles, involve one or more hosts. Hafeezullah, (1993) and Jadhav *et al.*, (2010) reported, host belongs to family Bagridae, Heteropneustidae, Schilbeidae and Siluridae, Mastacembelidae, Clariidae have been reported as definitive hosts of cestodes.

Parasitic infestation tends to decrease the growth rate resulting in stunting of the fish. The damage caused by helminths to their hosts is generally related to the intensity of infestation and the depth of parasite penetration with the host tissue. Seasonal fluctuation, locality, age, size and sex of the host also determine the parasitic community diversity and burden. Dogiel *et al.*, (1961) stated that seasonal environmental changes of water such as temperature, pH and conductivity affect on the occurrence of parasites from aquatic host. Parasitic diseases of fishes are very common all over the world. Globally the parasites (defined broadly as infectious agents of diseases) are responsible for 19% of human mortality (WHO, 2004). With the ever increasing demand for natural resources such as water due to high populations as well as minimizing the effects of drought, the government constructed dams which supply water to the local communities for farming, industry and general house hold use.

Such developments have brought with them benefits that can be exploited by the locals such as commercial and/or subsistence fishing activities. During the study of 'Population Dynamics and Seasonal Variations' of Platyhelminth parasites

MATERIALS AND METHODS

Sukhana dam is an earth fill dam on Sukhana River near Aurangabad Thasil. (Lay at 19°48'722"N and 75°31'096"E). Fishes like *Labeo rohita*, *Catla catla*, *Cirrhinus mrigala*, *Cyprinus carpio* and other exotic spp. like *Ctinopharyngdon*, *Hypothalamus* and *Chela* spp. are commonly found in this dam. Some of the carnivorous fishes such as *Ophiocephalus*, *Clarius*, *Mystus*, *Heteropneustes*, *Tilapia*, *Wallago*, etc. are commonly found in the reservoirs. The aim of the present study was to determine the helminth parasite fauna of fish, during the study period, an attempt was made to collect nearly an equal number of fishes of which most were of the small and medium size. The fishes were killed immediately before examination, since it was found that the small monogenetic trematodes rapidly disintegrated after the death of the fish. The examinations were made with a binocular stereoscopic microscope. When possible, helminths were removed from the host and identified while in the living state. Weight and length was measured after collection of fishes. Common necropsy and parasitological techniques were used to isolate the parasites. Parasites were preserved in 4% formalin. Worms were cleared in lactophenol, Parasites were preserved in 4% formalin, cleared in xylene, staining with hematoxylin, mounted in Canada balsam line drawings are made with the aid of drawing tube, identified by using standards keys (Khalil, 1991 and Parpena, 1996).

External examination on the gills, fins and surfaces of the fish for ectoparasites was first carried out using hand lens for detection of parasitic manifestations. Later, skin smear was made using scalpel blade. The procedure was performed using a spatula by which the skin scrapings (smears) from the head to the tail were obtained, mucus mixed with epidermal cells. Thereafter, the scraped samples of mucus together with the tissues were placed on a Petri-dish containing 3mls of 0.9% saline solution and stirred using a mounted pin (Omeji, et al., 2010, Bichi and Ibrahim, (2009) and Emere and Egbe, 2006). Some drops of the mixed solution were collected using dropper, placed on a clean slide and examined under microscope.

from Sukhana dam The prevalence, incidence, intensity, index of infection, abundance and dominance were calculated from collected cestode and trematode parasites along the graphical distribution.

The mean intensity was determined by dividing the total number of collected parasites by the number of infected fish samples, while abundance was calculated by dividing the total number of collected parasites by the number of host fish examined. The dominance of a parasite species was calculated as n/N sum (where n =abundance of a parasite species and N sum = sum of the abundance of all parasite species found).

RESULTS AND DISCUSSIONS

A total of 19 species of *Senga sukhaensis* were found in *Mastacembalus armatus* at Sukhana Dam; (73.68% of fishes are infected from Sukhana dam). Maximum intensity of infection (1.33) was found in February 2011 from Sukhana dam. 14 species of genus *Gangesia attuensis* were collected from *Wallago attu*, the incidence of infection 54.55% was found. Maximum abundance was recorded in the month of October 2011 at Sukhana dam. 10 species of *Phyllodistomum batrachusi* were collected from *Clarius batrachus* (63.33% of fishes are infected from Sukhana dam) high intensity (2.00) was recorded in the month of April 2012; one acanthocephala (*Acanthogyrus*) were found in *Clarias batrachus*, *Heteropneustes fossilis*, *Tilapia mossambica*, and *Wallago attu*. The fish biometrics is shown in Table 1 and Tables 2 shows the prevalence, mean intensity, range, abundance and dominance of the collected parasites.

14 species of genus *Gangesia attuensis* were collected from *Wallago attu* out of 22 host species, 12 fishes were found infected (54.55%), nematode, *Eustrongylides*, the acanthocephalan *Acanthogyrus* and piscine coccidian that was found in the intestines of host fishes. 19 species of *Senga sukhanaensis* were collected from *Mastacembalus armatus*, out of 95 host species 70 fishes were found infected (73.68%), total of 10 species of genus *Phyllodistomum batrachusi* collected from *Clarius batrachus* out of 30 host species 19 fishes were found infected (63.33%). During the study we collected nematodes, *P. cyathopharynx* from *Clarias batrachus*. The ectoparasite *Argulus* was collected from all fish species. Most *Argulus* was collected from the mouth cavity, gills and the fins.

Table 1 Biometric characteristic (weight and total length) of *Clarias batrachus*, *Wallago attu*, *Mastecembalus armatus* and *Clarius batrachus*

Fish	Weight (g) ±SD	Total length (cm) ±SD
<i>Mastecembalus armatus</i>	312.0±34.5 (200-428)	26.2±1.5 (23.2-28.4)
<i>Wallago attu</i>	306.0±33.5 (200-456)	28.8±1.6 (21.5-27.5)
<i>Clarias batrachus</i>	1078.9±187.3 (506-1220)	48.3±6.4 (31.5-51)

Table 2 The prevalence, mean intensity, abundance and dominance of Platyhelminth Parasites in freshwater fishes from Sukhana Dam during January 2010 to December 2012

Host	Parasites	No. of Infected fish	Prevalence %	Mean intensity ±SD	Range	Abundance ±SD	Dominance %
<i>M. armatus</i>	<i>S. sukhaensis</i>	70	73.68	0.27±0.06	0-8	0.20±0.05	0.006
<i>W. attu</i>	<i>G. attuensis</i>	12	54.55	1.17±0.01	0-6	0.64±0.05	0.020
<i>C. batrachus</i>	<i>P. batrachusi</i>	19	63.33	0.53±0.05	0-7	0.33±0.05	0.030

The current study showed that *Clarias batrachus* and *Mastacembalus armatus* had the greatest diversity of helminth parasites. Tapeworms are widespread throughout all over world (Campbell, 1999). Major water systems of India and demonstrate a high degree of host specificity, with Siluriform fish being the most common hosts for both monozoic and segmented cestodes. This could be attributed to the habitat favoured by *Clarias batrachus* that consists of turbid environments and shore areas which are covered with vegetation, as was the case of Jaikwadi Dam (Nimbalkar et al., 2010). This habitat also favours the intermediate hosts of cestodes as well as trematode digeneans. Hoffman in (1967) reported that in the mud habitat second intermediate hosts of many fish digeneans such as larvae of aquatic insects like Ephemeroptera, Odonata, Chironomidae and various Crustacea are found and form part of the diet of *Clarias batrachus*. Another reason for the recovery of a large number of helminths in *Clarias batrachus* could be related to the large size of *Clarias batrachus* as compared to other fish species (Table 1). The occurrence of very few helminth parasites in *Heteropneustes fossilis* and *Wallago attu* could be attributed to resistance to helminth infections.

Aquatic birds are important in the ecology of fish parasites because most helminths complete their life cycles in the bird host. Another interesting finding of our study was the recovery of piscine coccidia in *Clarias batrachus* only. *Coccidiosis* in fish usually manifests itself as a chronic infection and mortality

is gradual and overlooked in most fish farms. In conclusion, the obtained results show that helminths are important parasites of fishes in Sukhana dam, and detailed studies on the seasonal variations of these helminth parasites is recommended.

REFERENCE

Bichi A H and Ibrahim AA, 2009. A survey of ecto and intestinal parasites of tilapia zillii (gervias) in tiga lake, kano, Northern Nigeria. *Bayero Journal of Pure and Applied Sciences*, 2(1):79-82

Campbell NA, JB Reece and LG Mitchell, 1999. Biology. Benjamin Cummings.

Dogiel VA, Petrushevski GK, Polyanski YI, 1961. *Parasitology of fishes*. Leningrad: University Press; PMID: 13723441.

Emere MC and Egbe NEL, 2006. Protozoan Parasites of *Synodontis clarias* (A freshwater fish). *Best Journal*, 3(3):58-64.

Froese R and D Pauly, 2015. Fishbase. www.fishbase.org version (04/2015).

Hafeezullah M, 1993. Caryophyllidean cestode fauna of India. *Records of the Zoological Survey of India*, Occasional paper no. 157, 101.

Hoffman, 1967. Parasites of North American Fresh Water Fish University of California Press, Berkeley.

Jadhav BV, 2010. Survey of tapeworms from Aurangabad region. *Records of the Zoological Survey of India: A Journal of Indian Zoology*, 110 (Part-1): 107-114.

- Khalil LF, 1991.** Techniques for identification and investigative helminthology: Techniques for processing platyhelminths and acanthocephalans, in *Helminthology manual*, edited by L.F. Khalil. London: International Institute of Parasitology.
- Mittermeier RA, Mast RB, del Prado CP and Mittermeier CG, 1997.** In: R. A. Mittermeier, P. Robles Gil, and C.G. Mittermeier (eds.) *Megadiversity: Earth's Biologically Wealthiest Nations*. Monterrey, Mexico: CEMEX, 282-297.
- Nimbalkar RK, Shinde SS, Tawar DS and Nale VB, 2010.** A survey on Helminth Parasites of Fishes from Jaikawadi Dam, Maharashtra State of India. *Journal of Ecobiotechnology*. 2:8, 38-41
- Omeji S, Solomon SG and Obande RA, 2010.** A comparative study of the common protozoan parasites of *Heterobranchus longifilis* from the wild and pond environments in Benue State. *Pakistan Journal of Nutrition*. 9 (9): 865-872.
- Parpena I, 1996.** Parasites, infections and diseases of fishes in Africa: An update. *FAO/CIFA Technical Paper*, 31:130-199.
- Shukla GS and Upadhyay VB, 1998.** *A textbook of economic zoology*. Rastogi Publications, India, 205 p.
- Schmidt GD and LS Roberts, 2000.** Foundations of Parasitology. 6th edition Mc Graw-Hill International Editions, Boston. *Technical Paper*, 31:130-199.
- WHO, 2004.** Report of Joint WHO/FAO workshop on Food-borne trematodes infections in Asia, Ha Noi, Vietnam, 26-28 November, 2002. *World Health Organization*. WRPO, 1-58.

How to Cite this Article:

Nimbalkar R K and Deolalikar RV, 2015. Studies on Prevalence of Platyhelminth Parasites of freshwater fishes from Sukhana Dam at Aurangabad District (M.S.) India. *Bioscience Discovery*, 6(2):125-128.