



**Full Length Article**

## Four New Flowering plant Records from Satpuda Range of Jalgaon District, (MS) India

Tanveer A Khan<sup>1</sup>, Vivek V Desai<sup>2</sup> and N R Gawande<sup>3</sup>

<sup>1</sup>Department of Botany, H. J. Thim College of Arts and Science Mehrun, Jalgaon (M.S.)

<sup>2</sup>Cohort for Bio-Research Organization Jalgaon (M.S.)

<sup>3</sup>Range Forest Officer, Jamnya Range, Wildlife Sanctuary, Yawal (M.S.)

tanveerkhan04@gmail.com

### ABSTRACT

Jalgaon is one of the districts in the Khandesh region of North Maharashtra India with great biodiversity of plants. The present paper deals with the addition of 4 taxa of flowering plants to the flora of Jalgaon districts, Maharashtra. These taxa are *Habenaria plantaginea* Lindl. (Orchidaceae), *Euphorbiaserpens* Kunth. (Euphorbiaceae), *Exacum tetragonum* Roxb. (Gentianaceae), *Ceropegia hirsuta* Wight & Arn. (Asclepiadaceae). These species has been reported for the first time from Satpuda range of Jalgaon district, Maharashtra. The study provides a detailed taxonomic description, photographs and relevant information based on fresh collections.

**Key Words:** New records, Jalgaon districts, Satpuda ranges, Maharashtra.

### INTRODUCTION

The exploration of vegetation wealth of a region gives us correct understanding of bio-resources for the betterment of human beings. Jalgaon, Dhule and Nandurbar districts comprise Khandesh region, a northern part of Maharashtra. Jalgaon district lies between 20<sup>o</sup> and 21<sup>o</sup> North latitude and 74<sup>o</sup> 55' and 76<sup>o</sup> 28' East longitudes. The Jalgaon district has a total area about 272 sq. km. The total forest area in the district is 72685.27 hectares. On its location in the upper Tapi basin, it forms a distinct topographical unit separated from neighboring Madhya Pradesh state by Satpura ranges and from the south by Satmala hilly ranges. The physiographic of the district is made up of high hill ranges on the north, alluvium in the centre and low hill ranges to the south of Tapi. On the north, the hill ranges stretch east-west and form part of the Satpuras, the highest peak being about 1175 meters. The collected species like *Habenaria plantaginea* Lindl., *Euphorbia serpens* Kunth,

*Exacum tetragonum* Roxb., *Ceropegia hirsuta* Wight & Arn., are rare to Maharashtra and collected from only few places like Chandrapur, Bhandara, Raigad, Pune, Satara, Kolhapur, Ratnagiri, Singhudurg and Thane. The study region though botanically rich in biodiversity have not been explored extensively except a few sporadic reports on floristic of Khan 2014; More 2013; Kshirsagar 2008; Valvi 2006. The forest of Jalgaon district is of the tropical, dry deciduous type. The vegetation varies with the changes in altitude, aspect and rainfall. While working on floristic of Jalgaon district of Maharashtra we undertook frequent collection tours in every season to study plants. Many workers under botanical explorations in the Satpuda ranges (The range rises in eastern Gujarat state, running east through the border of Maharashtra and Madhya Pradesh to the east till Chhattisgarh) worked out many further additions to the flora of Satpuda, viz. Ray and Sainkhediya (2014), Kamble and Chaturvedi (2014).

**MATERIALS AND METHODS**

Satpuda ranges, which is one of the major hotspot of plants in Jalgaon district. During botanical exploration of Jalgaon district in Maharashtra four interesting species *Habenaria plantaginea* Lindl. (Orchidaceae) N 21°22.631' E 75°35.828' from Vaki, *Euphorbiaserpens* Kunth. (Euphorbiaceae) N 21°22.973' E 75°30.475' from Devgiri, *Exacum tetragonum* Roxb. (Gentianaceae) N 21°21.874' E 75°42.456' from Jamnya, while *Ceropegia hirsuta* Wight & Arn. (Asclepiadaceae) N 21°17.796' E 75°35.606' from Vagjira was collected from hill slopes, margins of water courses and tuff or limestone bedrock. The species was identified with the help of pertinent literature (Lakshminarasimhan *et al.*, 1996; Mitra 1971; Singh *et al.*, 2001; Mudgal *et al.*, 1997; Sardesai 2002) and the taxa were confirmed by Dr. Milind Sardesai Department of Botany, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad and by consulting the BSI western Circle, Pune, herbarium as well. The voucher specimens have been deposited in the herbarium of Department of Botany, H. J. Thim College of Arts and Science Mehrun, Jalgaon, Maharashtra.

**RESULTS AND DISCUSSION**

*Habenaria plantaginea* Lindl. Gen. Sp. Orchid. 323. 1835; Hook.f. Fl. Brit. India 6: 141. 1890; Sant. & Kapad. Orch. Bomba 28, t. 6, f. 25. 1966; Lakshmi. in Sharma *et al.* Fl. Maharashtra St. Monocot. 42. 1996. Sighet *et al.*, in Fl. M. P. 3: 45. 2001. (Plate I)

Terrestrial herbs, 20-50 cm high. Tubers 2, unequal. Leaves radical, prostrate on the ground, uppermost small and bract-like, oblong-lanceolate or elliptic, 4-8 x 1.5-2 cm, entire, membranous. Flowers in lax, few to many flowered racemes on long, slender scape, shortly pedicellate; bracts ovate-lanceolate, much shorter than ovary. Sepals white, unequal; dorsal sepal broadly ovate, entire; lateral sepals falcately oblong. Petals white, linear-lanceolate. Labellum rhomboid, 3-lobed, long spurred; mid lobe narrowly linear; lateral lobes much broader than mid lobe, wing like; spur white, slender, curved, pendulous, longer than ovary. Capsules fusiform, turgid, curved.

**Flowering and Fruiting:** September-November

**GPS Reading:** N 21°22.631' E 75°35.828' (Elevation 665<sup>m</sup>)

**Distribution:** Rare. In Satpuda ranges. In moist shady places in forest at high elevations. In

Maharashtra reported from Ahmednagar, Chandrapur, Bhandara, Raigad, Pune and Satara.

**Specimens examined:** Jalgaon Dist., Langdha Aamba, TAK 2398; Vaki, TAK 2472; Jamnya, TAK2587.

*Euphorbiaserpens* Kunth in Humb., Bonpl. & Kunth, Nov. Gen. Sp. 2: 52. 1817; Mitra in J. Bombay Nat. Hist. Soc. 68:854.1971; Mudgal *et al.*, in Fl. M. P. 2: 555. 1997. *Anisophyllum serpens* (Kunth) Klotzsch and Garcke, Abhandl. Akad. Berl. Phys. 1859: 23. 1860. *Chamaesyce serpens* (Kunth) Small, Fl. S.E.U.S. 709. 1903. *Euphorbia microphylla* auct. non Heyne ex Roth, 1821: Hook. f., Fl. Brit. India 5: 252. 1887, *p.p.* *Euphorbia orbiculata* Kunth var. *jawaharii* Rajagopal and Panigrahi in Taxon 17: 547. 1968. 'Matted Sandmat, Creeping Spurge' (Plate I).

A completely glabrous much-branched prostrate annual herb with stems to c. 25 cm long, rooting at the nodes. Petioles 0.5-1 mm long. Leaf-blades suborbicular-ovate, 1-5 x 1-4.5 mm, rounded to emarginated at the apex, obliquely shallowly cordate at the base, entire, pale green. Stipules interpetiolar, fused above and beneath to form a triangular lacinate or fimbriate white scale 0.5 mm long. Cyathia axillary, solitary, in the uppermost axils; glands transversely ovate or oblong, purplish, with narrow, subentire white appendages. Frutis trigonous, keels carinate, 1.2 x 1.5 mm, smooth. Seeds ovoid, quadrangular, 0.8 x 0.5 mm, smooth, pinkish-brown.

**Flowering and Fruiting:** August-November

**GPS Reading:** N 21°22.973' E 75°30.475' (Elevation 435<sup>m</sup>)

**Distribution:** Occasional. Along the banks of streams, canals and moist hill slopes.

**Specimens examined:** Jalgaon Dist., Manudevi forest, TAK 2235; Devgiri TAK2469; NMU Campus, TAK 2518.

**Note:** It looks similar to *E. heyneana* Spreng. but differs by its roots at nodes, green stem, sometimes with red striations, stipules on sides united, triangular, leaf blade oblong to sub-orbicular, leaf base truncate or cordate, white appendages of glands longer and wider than glands.

*Exacum tetragonum* Roxb. Fl. Ind. ed. Carey & Wall. 1: 413. 1820; Mudgal *et al.*, in Fl. M. P. 2: 104. 1997; Singh *et al.* Fl. Maharashtra St. Dicot. 2: 408. 2001. *E. tetragonum* var. *stylosa* (Wall. ex D. Don) Cl. in Hook. f. op. cit. *E. bicolor* Roxb. op. cit. 413; Cooke, Fl. Pres. Bombay 2: 252. 1958. (Repr.). 'Udi-chirayat'. (Plate I)

*Habenaria plantaginea* Lindl.*Euphorbia serpens* Kunth.*Ceropegia hirsuta* Wight & Arn.*Exacum tetragonum* Roxb.

### Plate I

Herbs, erect *ca* 60 cm high; stems 4-angled, branched in upper portion. Leaves lanceolate, 4-15 x 1-5 cm, narrowed at base or sessile. Flowers in terminal, much branched, paniculate cyme. Calyx lobes ovate, 5-10 mm long, acuminate, winged on the back. Corolla white with violet-blue patches towards edges; lobes lanceolate, 10-20 mm long, acute. Capsules globose, 0.7-1.5 cm long, smooth, shining, tipped with remains of style, yellowish-brown.

**Flowering and Fruiting:** September-February

**GPS Reading:** N 21°21.874' E 75°42.456' (Elevation 775<sup>m</sup>)

**Distribution:** Occasional, along the margins of streams, rice fields, among grasses along forest roads and in marshes of wet grasslands at high elevations. In Maharashtra reported from Kolhapur, Nasik, Raigad, Ratnagiri, Singhadurg and Thane.

**Specimens examined:** Jalgaon Dist., Jamnya TAK2352; Pathi TAK2536; Devgiri, TAK 2487.

**Note:** Leaves upper ones sessile, lower most sessile; corolla lobes more than 1.5 cm long.

***Ceropegia hirsuta*** Wight & Arn. in Wight, Contrib. 30. 1834; Singh *et al.* Fl. Maharashtra St. Dicot. 2: 350. 2001. Mudgal *et al.*, in Fl. M. P. 2: 69. 1997. 'Bosia kand, Hamana'. (Plate I)

Twining herbs, hispid; roots tuberous. Leaves variable, ovate, elliptic-lanceolate, 5-7 x 3-4 cm, acute to obtuse at apex, rounded at base, pilose. Flowers in few-flowered cymes; peduncles 1.2-2.5 cm long, stout, hispid. Corolla light green and variously striped with purple; lobes broad and folded back, upper 1/3<sup>rd</sup> light yellow, ciliate on margin and midrib. Outer corona-lobes deltoid, cleft at apex, pilose, extending above the gynostegium; inner corona-lobes hooked at tip, 3 times the length of the outer. Follicles up to 9 cm long.

**Flowering and Fruiting:** August-October

**GPS Reading:** N 21°17.796' E 75°35.606' (Elevation 422<sup>m</sup>)

**Distribution:** Occasional. Along the hill slopes and in moist rocky places. Elsewhere in Maharashtra reported from Akola, Aurangabad, Nasik, Pune, Ratnagiri and Satara.

**Specimens examined:** Jalgaon Dist., Vagjira, TAK2592; Mandap-Nalah, TAK2609; Vaki, TAK 2479.

**Note:** Plants hispid; corolla tube hairy inside; inner corona hooked at tips.

**Uses:** The decoction of plant is given orally to antidote snake bite coupled with the external application of plant paste (Mudgal 1997).

**CONCLUSION**

We have gone through all pertinent literature (Kshirsagar 2008, Patil 2003) and by consulting the BSI Herbarium Pune. To find out the occurrence, distribution and habitat of these species. We found that, these species were not reported in any of the Jalgaon flora. This clearly reveals that, these species are rare to flora of Maharashtra State, even India as a whole. These species are new record to the flora of Jalgaon district of Maharashtra State. The voucher specimens are deposited in the herbarium of Department of Botany, H. J. Thim College of Arts and Science Mehrun, Jalgaon.

On close examination of herbarium specimens and detailed scrutiny of literature published till today on these taxa, it can be claimed that these are new records for Satpuda range of Jalgaon district of Maharashtra.

**ACKNOWLEDGEMENT**

The authors wish to express their gratitude to Dr. Milind Sardesai, (Department of Botany, Dr. Babasaheb Ambedkar Marathwada University, Aurangabad) who confirmed the identity of these species. One of the author (TAK) is thankful to North Maharashtra University for financial support under the (VCRMS). Thanks are also due to the Principal, H.J. Thim College, Jalgaon, for providing laboratory and library facilities.

**REFERENCES**

**Hook JD, 1890.** *Flora of British India* L. Reeve and Co. London, **6**:140-143.  
**Kamble RB and Chaturvedi A, 2014.** New additions to the Flora of Nagpur District, Maharashtra – IV. *Biosciences Discovery*, **5**(2):160-162.

**Khan TA and Chaudhari GS, 2014.** New records of Cyperaceae for Khandesh region of Maharashtra, India. *Plant Archives*, **14**(1): 235-240

**Kshirsagar SR and Patil DA, 2008.** *Flora of Jalgaon district, Maharashtra*. Bishen Singh Mahendra Pal Singh, Dehradun, 41-376.

**Lakshminarasimhan P, Sharma BD, Karthikeyan S and Singh NP, 1996.** *Flora of Maharashtra State. (Monocotyledons)* Botanical survey of India, Kolkata, 42-43.

**Mitra RL, 1971.** *Euphorbia serpens* H. B. K. (Euphorbiaceae) a hitherto unrecognised species in India. *Journal Bombay Natural Historical Society*, **68** (3): 852-856.

**More NK, Kamble SS and Dhabe AS, 2013.** Some new records for the flora of jalgaon district, (m.s.) India. *Bioinfolet*, **10**(4a): 1108-1109.

**Mudgal V, Khanna KK, Hajara PK, 1997.** *Flora of Madhya Pradesh*, Botanical Survey of India, Kolkata, **2**: 69-555.

**Patil DA, 2003.** *Flora of Dhule and Nadurbar District (Maharashtra)* Bishan Singh Mahendra Pal Singh Deharadun, 53-649.

**Rajagopal T and Panigrahi G, 1968.** The correct name for *Euphorbia microphylla* and a new variety, *J. Taxon* **17**: 547.

**Ray S. and Sainkhediya J, 2014.** Some New Record for the flora of Madhya Pradesh. *Bioscience Discovery*, **5**(2):187-192.

**Singh NP, Khanna KK, Mudgal V and Dixit RD, 2001.** *Flora of Madhya Pradesh*, Botanical Survey of India, Kolkata, **3**: 45-46.

**Singh NP, Lakshminarasimhan P, Karthikeyan S and Prasanna PV, 2001.** *Fl. Maharashtra St. Dicot.*, **2**: 350-409. 2001.

**Valvi RJ, Yadav SS, and Varghese M, 2006.** New record of orchid species for the flora of West Khandesh Satpuda. *Plant archives*, **6**(2): 753-755.

How to Cite this Article:

**Tanveer A Khan, Vivek V Desai and N R Gawande, 2015.** Four New Flowering plant Records from Satpuda Range of Jalgaon District, (MS) India. *Bioscience Discovery*, **6**(1):45-48.