



Mercury toxicity; behavioral changes and recovery in a freshwater teleost *Heteropneustes fossilis* (Bloch)

Bhalerao Sangeeta N

Department of Zoology,
Anantrao Pawar College of Arts, Commerce and Science, Pirangut, Tal. Mulshi, Pune (MS) India
bhalerao.sangeeta@rediffmail.com

Article Info

Received: 11-11-2016,
Revised: 26-11-2016,
Accepted: 30-12-2016

Keywords:

Mercury, Recovery,
Toxicity, Heavy Metal, *H. fossilis*, Pollution.

Abstract

Present study was aimed to observe the behavioral changes due to Hg (0.1mg/l) toxicity in the freshwater catfish *Heteropneustes fossilis* (Bloch). The fishes exhibited erratic swimming, thick mucus secretion over body surface, skin soars and jerky movements. Fishes tried to leap out of aquarium during experiments. It was further observed that these behavioral changes were suppressed when fishes were simultaneously fed with herbal compound. The recovery experiments confirmed that the behavioral changes due to Hg toxicity were reduced when fed with the herbal compound, indicating its protective effects against Hg toxicity.

INTRODUCTION

The aquatic life of water resources is in danger due to industrial pollutants, domestic, agricultural wastes, heavy metals, etc. Numerous studies confirmed that pollutants are adversely affecting the aquaculture (Opaluwa *et al.*, 2012; Haloi *et al.*, 2013). The presence of pesticides and various toxic metal accumulations were observed and reported in the freshwater and marine fish organs (Ali and Fishar, 2005; Agrahari S and Gopal K, 2007; Karthigayani *et al.*, 2014). It is well-known that Mercury is a highly toxic non-essential heavy metal. In the modern age its application is increasing in industries and agriculture. It gets accumulated and damages the fish organs. The higher concentrations of Hg damage the organs and intern the fish behavior. Consumption of such fishes through the natural food chain reaches to various human tissues (Devlin, 2006).

Any changes in fish behavior are considered as one of the sensitive biomarker to evaluate the exposure to the toxicant (Gerhardt,

2007; Reddy *et al.*, 2011). It had been observed that the fish behavior alters due to the toxicants like heavy metals, pesticides, etc. (Javed, 2012; Ghanbahadur, *et al.*, 2015 and Deshmukh, 2016). Erratic swimming, escaping from toxic water, mucus secretion, convulsions and food intake, etc. can be considered as some common parameters to measure the changes in fish behavior. The preventive and curative effects were also studied by researchers. The herbal compound Liv₅₂ is a hepatoprotective drug (Rathore and Verma, 1988; Jain, 2016) and was examined against Hg and other toxicants in fresh water fishes (Kumar and Kothari, 1990, Palas *et al.*, 2014) and in mammals (Keshav and Lal, 2016).

In paucity of literature of the protective action of herbal compound Liv₅₂ against Hg toxicity and behavioral changes in fresh water teleost *H. fossilis*, the present study was aimed. The role of herbal compound Liv₅₂, if any, in the recovery process of the Hg exposed fish was also examined.

MATERIALS AND METHODS

The fresh water fish *H. fossilis* was procured from the local water body of Ujjain (MP). The fishes were acclimatized to laboratory conditions in glass aquaria for seven days. Stock solution of HgCl₂ was prepared as per the guidelines of APHA (1975). The experimental concentration of HgCl₂ was kept at 0.1 mg/l. The indigenous hepatoprotective herbal compound Liv₅₂ was tested for its protective role. The food and drug were fed at the rate of 30 and 10 mg/day/fish, with few drops of liquid paraffin, to the fishes of all aquaria respectively. On every fourth day water of all aquaria was changed and fresh metal solution was added to experimental groups.

The acclimatized fishes were divided into groups of 25 each as under:

Group I - Control fed on normal food

Group II - Treated with HgCl₂ (0.1 mg /l) and fed on normal food

Group III - Treated with HgCl₂ (0.1 mg/l) and fed on food containing drug (Hg + drug)

Group IV - First 30 days treated with HgCl₂ (0.1 mg/l) and fed on normal

food then divided into two groups

Group IV A - 30 days kept in Hg free water and fed on normal food, i.e., group IV A, Natural recovery

Group IV B - 30 days kept in Hg free water and treated with drug, i.e., group IV B Drug recovery

The fishes of these groups were observed daily. The behavioral changes in all the groups were compared with the control group fishes. Fishes from groups I, II and III were observed for 30 days, whereas the fishes of recovery groups were observed for 60 days.

RESULTS AND DISCUSSION

The fresh water fish *H. fossilis* exposed to HgCl₂ showed changes in its normal behavior. The fishes of group II, exhibited a change in swimming pattern. During initial exposure, fishes showed jerky movements with rapid and erratic swimming. They exhibited tendency of escaping from the aquarium. On prolonged Mercury exposure the fishes became sluggish and showed uncoordinated swimming patterns. The similar results were observed in recent past (Sabullah *et al.*, 2015; Sehar *et al.* 2014). This may be due to the toxic Hg interference in some of the functions which were controlled by the nervous system of the fishes (Daoust 1981; Thangam and

Manju, 2013). In group III (Hg + drug) the jerky movements and erratic swimming were reduced.

An excessive mucus secretion over the body surface was observed in the prolonged Hg exposed fishes. This can be attributed to the increased stress due to toxic effect of Hg and caused changes in the area and number of mucus glands (Akhter and Noori, 2014). The excessive secretion of mucus changed the colour of aquaria water to milky white. During drug recovery (group IV B), experiments fishes exhibited normal swimming behavior and no excessive mucus secretion.

Like other toxicants the mercury toxicity also caused reduction in food intake in the fishes. The food was not fully consumed in metal exposed fishes as against the control group. It was observed that the food intake was slightly higher in group III (Hg + Liv₅₂) as compared to the group II (Hg exposed) fishes. The food intake of the recovery group (IV B) fishes was comparable with the control group fishes. This suggested that Hg caused loss of appetite and the Liv₅₂ helped to maintain the normal food intake. Liv₅₂ had been reported to increase food consumption in laboratory animals (Reddy *et al.*, 1990; Palas *et al.*, 2014). The efficacy of hepatoprotective herbal compound Liv₅₂ with *Ricinus communis* in Wistar rats was also reported in recent past (Navin *et al.*, 2016).

The findings of this investigation revealed that the intoxication of Hg caused uncoordinated swimming activity, excess mucus secretion and also decreased food consumption in *H. fossilis*. This may be due to the damage to nervous system by Hg intoxication, whereas, Liv₅₂ helped to maintain normal behavior, especially food intake. It can be concluded that the herbal compound (Liv₅₂) was almost found effective in controlling the Hg induced abnormal behavior. However, further detailed studies are needed to understand the mode of protective action of the herbal compound (Liv₅₂) against toxic action of Hg.

REFERENCES

- Agrahari S and Gopal K, 2007.** Fate and toxicity of Cadmium and Lead accumulation in different tissues (gills, liver, kidney, brain) of fresh water fish, *Channa punctatus*. *J. Ecophysiol. Occup. Hlth.* **7**: 151-155.
- Akhter AS and Noori A, 2011.** Toxicity of heavy metal copper and its effect on the behaviour of fresh water Indian cat fish, *Clarias batrachus* (Linn.). *CB.*, **4**(4), 2011

- Ali MH and Fishar MR, 2005.** Accumulation of trace metals in some benthic invertebrate and fish species relevant to their concentration in water and sediment of lake Quaran, Egypt. *J. Aquat. Res.*, **31**: 289-301.
- APHA AWWA and WPCF, 1975.** Standard methods for the examination of water and waste water. 14th Edition, *American Public Health Association, Washington*.
- Daoust PY, 1981.** Acute pathological effects of Mercury, Cadmium and in rainbow trout. PhD Thesis, *University of Saskatchewan, Canada*. 331 pp.
- Deshmukh DR, 2016.** Hamatological response in a freshwater fish *Channa striatus* exposed to endosulfan pesticide. *Bioscience Discovery*, **7** (1):67-39.
- Devlin EW, 2006.** Acute toxicity, uptake and histopathology of aqueous methyl mercury to fathead minnow embryos. *Ecotoxicol.* **15**: 97-110.
- Gerhardt A, 2007.** Aquatic behavioral ecotoxicology – prospects and limitations. *Human and Ecological Risk Assessment.* **13**: 481-491.
- Ghanbahadur AG, Ghanbahadur GR, Ganeshwade RM and SB Wagh, 2015.** Histopathological effect of organochloride endosulfan on gills of larvivorous fish *Rarbora daniconius*. *Bioscience Discovery*, **6** (2):121-124.
- Haloi K, Kalita M and Nath R, 2013.** The study on the histopathological changes of stomach of *Channa punctatus* (Bloch). By used pesticide endosulfan. *Global J Sci Front Res Biol Sci* **13**.
- Jain N, 2016.** Western Himalaya-a treasure of pharmacological useful drugs. *iP-Planet*, **4** (1), 36-40, 2016.
- Javed M, 2012.** Growth responses of fish under chronic exposure of waterborne and dietary metals. *International Journal of Agriculture.* **14**: 281-285.
- Karthigayani T, Denis M, Andrew RAR and Shettu N, 2014.** Histological study of the intestine and liver tissues in the fish *Oreochromis mossambicus* exposed to cypermethrin. *Journal of Modern Biotechnology.* **3**:35-41.
- Keshav A and Lal UR, 2016.** Hepatitis and medicinal plants: An overview. *Journal of Pharmacognosy and Phytochemistry.* **5**(6): 408-415.
- Kumar R and Kothari SC, 1990.** Sites of Mercury accumulation in the Liver of *H. fossilis*, correlation between the structural damage and sites of accumulation. In: *Environmental pollution and Health Hazards Environmental series* vol. **III** (Eds. R. Prakash and S. N. Choubey) 127-132.
- Naveen A, Shankar J, Premendran J and Venkatanarayana N, 2016.** Evaluation of hepatoprotective activity of aqueous extract of *Ricinus communis* in Wistar rats. *Int J Basic Clin Pharmacol.* **5** (2):358-361.
- Opaluwa OD, Aremu MO, Ogbo LO, Magaji JI, Odiba IE and Ekpo ER, 2012.** Assessment of Heavy Metals in Water, Fish and Sediments from UKE Stream, Nasarawa State, Nigeria. *Current World Environment.*, **7**(2), 213-220.
- Palas S, Pal S, MukharjeeAK and Ghosh AR, 2014.** Evaluation of Metabolic Enzymes in Response to Excel Mera 71, a Glyphosate-Based Herbicide, and Recovery Pattern in Freshwater Teleostean. *Fishes.BioMed Research International.*, Volume Article ID **425159**, 6 pages.
- Rathore HS and Verma R, 1988.** Protection of mice liver with Liv₅₂ against cadmium intoxication. *Probe* **28**: 8-16.
- Reddy JS, Reddy TK and Reddy DC, 2011.** Influence of Heavy metals on biochemical and metabolic biomarkers of Indian Major Carp, *Labeo rohita*. *The Bioscan.* **6**(1): 167-173.
- Reddy PB, Kothari SC and Rathore HS, 1990.** Protective role of Liv₅₂ against histological damage due to CdCl₂ toxicity in the intestine of a teleost fish, *Probe*, **29**, 220-228.
- Sabullah MK, Ahmad SA, Shukor MY, Gansau AJ, Syed MA Sulaiman MR and Shamaan NA, 2015.** Heavy Metal biomarker: Fish behavior, cellular alteration, enzymatic reaction and proteomics approaches. *IFRJ.* **22** (2): 435-454.
- Sehar A, Shafaqat A, Uzma SA, Mujahid F, Saima AB, Fakhir H and Rehan A, 2014.** Effect of different Heavy Metal Pollution on Fish. *Res J. Chem. Environ. Sci.*, **2** (2): 35-40.
- Thangam Y and Manju M, 2013.** Mercury Toxicity in Hormonal Effects to Fresh Water Fish *Cirrhinus Mrigala*. *IJSR*, Paper ID: **0310150**, 1669-1675.

How to Cite this Article:

Bhalerao Sangeeta N, 2017. Mercury toxicity; behavioral changes and recovery in a freshwater teleost *Heteropneustes fossilis* (Bloch). *Bioscience Discovery*, **8**(1):61-63.